



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## SOLUTION BY W. J. THOME, University of Detroit.

It is a well known fact that if the parallel sides of a trapezoid are  $g$  and  $s$ , and the perpendicular distance between them is  $p$ , then the distance from the greater side  $g$  to the center of gravity of the trapezoid is

$$\frac{p}{3} \frac{(g + 2s)}{(g + s)}.$$

Let  $t$  be the constant width of the lever, in inches;

$d$ , the density of the material of the lever,

$M$ , the total mass of the lever;

$m_1$ , the mass of that part of the lever from the knife edge to the end distant  $a$  feet away;

$m_2$ , the mass of that part of the lever from the knife edge to the end distant  $b$  feet away;

$\bar{X}$ , the distance in feet from the knife edge to the center of gravity of the entire lever;

$\bar{x}_1$ , the distance in feet from the knife edge to the center of gravity of the mass  $m_1$ ; and

$x_2$ , the distance in feet from the knife edge to the center of gravity of the mass  $m_2$ .

Then, using the foot as our unit of length, we have

$$M\bar{X} = m_2\bar{x} - m_1\bar{x}_1;$$

or

$$\begin{aligned} & \left[ \frac{1}{2} \left( \frac{m+n}{12} \right) a \frac{t}{12} d + \frac{1}{2} \left( \frac{m+n}{12} \right) b \frac{t}{12} d \right] \bar{X} \\ &= \left[ \frac{1}{2} \left( \frac{m+n}{12} \right) b \frac{t}{12} d \right] \left[ \frac{b \left( \frac{m+2n}{12} \right)}{3 \left( \frac{m+n}{12} \right)} \right] - \left[ \frac{1}{2} \left( \frac{m+n}{12} \right) a \frac{t}{12} d \right] \left[ \frac{a \left( \frac{m+2n}{12} \right)}{3 \left( \frac{m+n}{12} \right)} \right]. \end{aligned}$$

Hence,

$$\bar{X} = \frac{1}{3} \frac{(m+2n)}{(m+n)} (b-a),$$

which is to be measured in the direction of whichever is the greater,  $b$  or  $a$ .

## NUMBER THEORY.

## 240. Proposed by J. W. NICHOLSON, Louisiana State University.

If the roots of  $x^3 - px + q = 0$  are rational, prove that  $4p - 3x^2$  is a perfect square.

SOLUTION BY WILLIAM E. PATTEN, Government Institute of Technology, Shanghai, China.

Let the roots of  $x^3 - px + q = 0$  be  $x_1, x_2, x_3$ . Then

$$x_1 + x_2 + x_3 = 0;$$

and

$$x_1x_2 + x_2x_3 + x_3x_1 = -p.$$

Therefore,

$$\begin{aligned} 4p - 3x_1^2 &= -4(x_1x_2 + x_2x_3 + x_3x_1) - 3x_1^2 \\ &= -4x_2x_3 - 4x_1(x_2 + x_3) - 3x_1^2 \\ &= -4x_2x_3 - 4(-x_2 - x_3)(x_2 + x_3) - 3(-x_2 - x_3)^2 \\ &= -4x_2x_3 + (x_2 + x_3)^2 = (x_2 - x_3)^2. \end{aligned}$$

Since the coefficient of the leading term of the given equation is unity, and the roots are rational, they are also integral.

Therefore,  $x_2 - x_3$  is an integer, and  $4p - 3x_1^2$  is a perfect square.

Similarly for the other roots:

$$4p - 3x_2^2 = (x_3 - x_1)^2, \quad \text{and} \quad 4p - 3x_3^2 = (x_1 - x_2)^2.$$

Also solved by H. N. CARLETON, J. E. ROWE, L. C. MATHEWSON, A. H. HOLMES, ELIJAH SWIFT, O. S. ADAMS, J. ROSENBAUM, LOUIS CLARK, E. E. WHITEFORD, and H. S. UHLER.

**241. Proposed by CLIFFORD N. MILLS, Brookings, S. Dak.**

If  $a^2 + b^2 = c^2$ , where  $a$ ,  $b$ , and  $c$  are integers, then prove that  $abc$  will be a multiple of 60.

SOLUTION BY ALBERT G. CARIS, Defiance College.

From the well-known theorem that any integral solution of  $a^2 + b^2 = c^2$  may be put in the form  $2xy$ ,  $x^2 - y^2$ ,  $x^2 + y^2$ , where  $x$  and  $y$  are integers, it follows immediately that

$$abc = 2xy(x - y)(x + y)(x^2 + y^2).$$

Showing that this product is always a multiple of 3, 4, and 5 is sufficient to prove the proposed problem.

I. We may write  $x = 2m - 1$ , or  $2m$  and  $y = 2n - 1$ , or  $2n$ , where  $m$  and  $n$  are integers. Whenever  $x = 2m$ , or  $y = 2n$ ,  $abc$  is a multiple of 4. In all other cases  $x = 2m - 1$  at the same time that  $y = 2n - 1$  and consequently,  $x - y$ ,  $x + y$ , and  $x^2 + y^2$  are all multiples of 2.

Therefore  $abc$  is always a multiple of 4.

II. We may write  $x = 3r - 1$ ,  $3r$ , or  $3r + 1$  and  $y = 3s - 1$ ,  $3s$ , or  $3s + 1$ , where  $r$  and  $s$  are integers. Whenever  $x = 3r$ , or  $y = 3s$ ,  $abc$  is a multiple of 3. The combinations resulting from all other cases may be arranged in the two groups below:

Group A		Group B	
$x$	$y$	$x$	$y$
$3r - 1$	$3s - 1$	$3r - 1$	$3s + 1$
$3r + 1$	$3s + 1$	$3r + 1$	$3s - 1$

From combinations of group A the factor  $x - y = 3(r - s)$ . From combinations of group B the factor  $x + y = 3(r + s)$ . Therefore  $abc$  is always a multiple of 3.

III. We may write

$$x = 5u - 2, \quad 5u - 1, \quad 5u, \quad 5u + 1, \quad \text{or} \quad 5u + 2$$

and

$$y = 5v - 2, \quad 5v - 1, \quad 5v, \quad 5v + 1, \quad \text{or} \quad 5v + 2,$$

where  $u$  and  $v$  are integers. Whenever  $x = 5u$ , or  $y = 5v$ ,  $abc$  is a multiple of 5. The combinations resulting from all other cases may be arranged in the three groups below:

Group C		Group D		Group E	
$x$	$y$	$x$	$y$	$x$	$y$
$5u - 2$	$5v - 2$	$5u - 2$	$5v + 2$	$5u \pm 2$	$5v \pm 1$
$5u - 1$	$5v - 1$	$5u - 1$	$5v + 1$		
$5u + 1$	$5v + 1$	$5u + 1$	$5v - 1$	$5u \pm 1$	$5v \pm 2$
$5u + 2$	$5v + 2$	$5u + 2$	$5v + 2$		

All combinations of group C make  $x - y = 5(u - v)$ . All combinations of group D make  $x + y = 5(u + v)$ . All combinations of group E make  $x^2 + y^2 = 5(5u^2 \pm 4u + 5v^2 \pm 2v + 1)$  or  $5(5u^2 \pm 2u + 5v^2 \pm 4v + 1)$ . Therefore,  $abc$  is always a multiple of 5. Hence,  $abc$  is always divisible by 60.

Also solved by S. A. COREY, A. H. HOLMES, HORACE OLSON, J. W. CLAWSON, J. ROSENBAUM, J. E. ROWE, H. C. FEEMSTER, H. N. CARLETON, W. J. THOME, ELIJAH SWIFT, and E. E. WHITEFORD.

**243. Proposed by CLIFFORD N. MILLS, Brookings, South Dakota.**

Determine rational values of  $x$  that will render  $x^3 + px^2 + qx + r$  a perfect cube. Apply the result to  $x^3 - 8x^2 + 12x - 6$ .